Tri-Permuting Mixing Matrix and predictions for θ_{13}

Federica Bazzocchi[1],

SISSA and INFN, Sezione di Trieste,
via Bonomea 265, 34136 Trieste, Italy

arXiv:1108.2497v1 [hep-ph] 11 Aug 2011

Abstract

We introduce a new texture for neutrino mixing named Tri-Permuting (TP) mixing matrix. This pattern is characterized by maximal solar and atmospheric angles and by a large reactor angle satisfying $\sin\theta_{13} = 1/3$. The correct lepton mixing matrix is obtained when combining the charged lepton mixing matrix with the neutrino one. In this way we get new predictions for θ_{13} with respect to those obtained by the well studied TBM or BM mixing patterns. We present a specific model that gives rise to TP mixing in the neutrino sector as well as the required corrections from the charged lepton one.

1 Introduction

The most recent T2K [1] and MINOS [2] results have shown at 2.5 and 1.7 σ respectively the evidence of a $\theta_{13} \neq 0$ in the lepton mixing matrix. The first global analysis [3] have confirmed their results giving at 3σ level the range $0.001 \leq \sin\theta_{13}^2 \leq 0.044$ ($0.005 \leq \sin\theta_{13}^2 \leq 0.050$) for the NH (IH) case. Comparable results have been obtained by the most recent global analysis [4] that have slightly lowered the upper bound $0.001 \leq \sin\theta_{13}^2 \leq 0.035$ ($0.015 \leq \sin\theta_{13}^2 \leq 0.039$) for the NH (IH) case.

While waiting for more statistics and forthcoming tests of these results neutrino phenomenology community have showed an impressive fast and conspicuous productivity in proposing new textures and models that could account for the correct θ_{13} size. The majority of these analysis have been devoted to a re-consideration of the possible corrections to TriBiMaximal (TBM) mixing. TBM pattern predicts at leading order (LO) a vanishing θ_{13}. Even in the first TBM predicting models [5] a non vanishing θ_{13} was indeed predicted at next to leading order (NLO) typically quite small, of order θ_C^2, with $\theta_C \sim .23$ the Cabibbo angle. In the last months different scenarios based mainly on discrete symmetries have been proposed to modify TBM texture and predicting a $\theta_{13} \neq 0$ up to 10 degrees [6, 7]. Other possibilities have been considered in [8].

Even before T2K and MINOS recent data a promising idea to get a $\theta_{13} \neq 0$ was given by BiMaximal (BM) mixing.

In the context of BM mixing the basic idea is that at LO solar and atmospheric angles are maximal and the reactor angle is zero [9–11]. Then at next leading order (NLO) only the solar and the reactor angles get corrections of order the Cabibbo angle θ_C while the atmospheric keeps unchanged. Finally at next-next leading order (NNLO) even the atmospheric angle may get corrections but these are of order θ_C^2 thus allowing to fall in the experimental data range. In this picture NLO corrections at the lepton mixing matrix arise by diagonalizing the charged lepton sector. The reason is seeded in the original motivation to study BM mixing, that is quark-lepton complementarity [12]. We remind that recently it has been shown how a relative large θ_{13} may arise from the charged lepton sector in the context of SU(5) assuming exact TBM mixing in the neutrino sector [7].

---

[1] e-mail address: fbazzo@sissa.it

At the moment while we are looking for more statistics and new analysis to confirm and deline the θ_{13} range one of the challenges is finding a texture that has a non vanishing θ_{13} at LO–eventually even too large–and thus making it smaller thanks to adequate corrections [13].

At the light of the most recent results there is an intrinsic tension in the BM assumption. Consider the BM choice

$$\theta_{12} = \theta_{23} = -\frac{\pi}{4}\,, \quad \theta_{13} = 0\,, \tag{1}$$

and assume that the charged lepton mass matrix is diagonalized on the left by a rotation in the 12 sector of order the Cabibbo angle parametrized as

$$U_e = \left(\begin{array}{ccc} \cos\theta & \sin\theta e^{i\delta} & 0 \\ -\sin\theta e^{i\delta} & \cos\theta & 0 \\ 0 & 0 & 1 \end{array} \right)\,, \tag{2}$$

where δ is a possible CP Dirac phase.

Then one finds that the lepton mixing angles are given by

$$\begin{array}{rcl} \theta_{23} & \sim & -\frac{\pi}{4} + \frac{1}{4}\theta^2 + \mathcal{O}(\theta^3)\,, \\ \theta_{12} & \sim & -\frac{\pi}{4} - \frac{1}{\sqrt{2}}\theta\cos\delta + \mathcal{O}(\theta^3)\,, \\ \theta_{13} & \sim & \frac{1}{\sqrt{2}}\theta\,. \end{array} \tag{3}$$

The solar angle wants a *large* θ of order the Cabibbo angle θ_C, while the most recent fit indicates that θ_{13} is large but not too much. The results is shown in fig. 1: according to the simple parametric expansion of eq. (3) the BM θ_{13} prediction is large and could be ruled out by an improvement of precision that could low the 3σ upper bound. However in more realistic scenarios the values allowed for θ_{13} are spreaded, but we still may conclude that if the upper 3σ limit on θ_{13} would be lowered the BM pattern would be strongly disfavored.

In this paper we present a complementary picture to that offered by the BM pattern that we define as the Tri-Permuting (TP) mixing matrix. The name reminds that the 3 eigenvectors are identical up to permutations and change of signs. This TP mixing matrix is defined by two maximal angles and a large θ_{13} according to

$$\sin\theta_{12} = \sin\theta_{23} = -\frac{1}{\sqrt{2}}\,, \quad \sin\theta_{13} = \frac{1}{3}\,. \tag{4}$$



Figure 1: *The reactor versus the solar angle in the case of the BM mixing matrix corrected by a rotation in the 12 plane of the charged lepton mass matrix. Vertical and horizontal lines bound the* 3σ *range for* $\sin\theta_{12}^2$ *and* $\sin\theta_{13}^2$ *respectively.*

$$U_{TP} \sim \frac{1}{3}\begin{pmatrix} 2 & -2 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 2 \end{pmatrix} . \tag{5}$$

Under the previous assumption that charged lepton mass matrix is diagonalized through the rotation given in eq. (2) we get

$$\begin{aligned} \theta_{23} &\sim -\frac{\pi}{4} + \frac{1}{4}\theta\cos\delta + \frac{1}{16}\theta^2(4+\cos 2\delta) + \mathcal{O}(\theta^3)\,, \\ \theta_{12} &\sim -\frac{\pi}{4} - \frac{3}{4}\theta\cos\delta - \frac{3}{16}\theta^2\cos 2\delta + \mathcal{O}(\theta^3)\,, \\ \theta_{13} &\sim \frac{1}{3} + \frac{2}{3}\theta\cos\delta - \frac{1}{6}\theta^2(1+4\sin\delta^2) + \mathcal{O}(\theta^3)\,. \end{aligned} \tag{6}$$

Notice that while θ_{12} and θ_{13} receive a correction of order $\sim \theta\cos\delta$, θ_{23} is corrected by $\theta\cos\delta/4 \sim \theta_C^2$ if $\theta \sim \theta_C$ the Cabibbo angle. Moreover to constrain θ_{12} in the correct range we need $\cos\delta < 0$ that gives a correction to θ_{13} in the right direction. This is explicitly shown in fig. 2. At the same time we get a prediction for the Dirac CP phase that in this scenario is given by

$$\delta_l \sim 2(\theta\sin\delta - \theta^2\sin 2\delta)\,. \tag{7}$$

In the next section we introduce the framework in which TP mixing matrix arises. In sec.3 we build a renormalizable model that provides such a texture. Neutrino phenomenological implication are discussed in sec.4 and sec.5 is devoted to our conclusions.



Figure 2: *The reactor versus the solar angle in the case of the TP mixing matrix corrected by a* 12 *rotation in the charged lepton sector . Vertical and horizontal lines bound the* 3σ *range for* $\sin\theta^2_{12}$ *and* $\sin\theta^2_{13}$ *respectively.*

2 Residual symmetries

It is well known that under the assumption that neutrinos are majorana particles if there is any residual symmetry behind neutrino mass matrix this is at most a $Z_2 \times Z_2$ flavor symmetry [13, 14]. While it is clear how the $Z_2 \times Z_2$ acts on the neutrino mass eigenstates since the three mass eigenstates must have *flavor parity* (+,+),(+,-),(-,-), it is an open question how it acts on the neutrino interaction eigenstates. In the most general case given the three left handed neutrinos $\nu_L \sim (\nu_{L_1}, \nu_{L_2}, \nu_{L_3})$ each Z_2 flavor symmetry $S_{1,2}$, $S^2_{1,2} = I$, acts on ν_L as

$$\nu_L \to S_i \nu_L \tag{8}$$

and it holds that $[S_1, S_2] = 0$. Thus the neutrino mass matrix may be written in terms of the three eigenvectors of S_1 and S_2, v_i, satisfying

$$\begin{aligned} S_1 v_1 &= v_1 & S_2 v_1 &= v_1\,, \\ S_1 v_2 &= v_2 & S_2 v_2 &= -v_2\,, \\ S_1 v_3 &= -v_3 & S_3 v_3 &= -v_3\,. \end{aligned} \tag{9}$$

As consequence the effective light neutrino majorana mass matrix may be written as

$$m_\nu = m_1 v_1^T v_1 + m_2 v_2^T v_2 + m_3 v_3^T v_3\,. \tag{10}$$

This approach has been used in many scenarios and typically addressed as sequential dominance [15]. Determining S_1 and S_2 fixes uniquely the lepton mixing matrix if the charged lepton mass matrix is diagonal. S_1 and S_2 corresponding to the TP mixing matrix are given by

$$S_1 = \left(\begin{array}{ccc} \frac{7}{9} & \frac{4}{9} & -\frac{4}{9} \\ \frac{4}{9} & \frac{1}{9} & \frac{8}{9} \\ -\frac{4}{9} & \frac{8}{9} & \frac{1}{9} \end{array} \right) , \qquad S_2 = \left(\begin{array}{ccc} -\frac{1}{9} & \frac{8}{9} & \frac{4}{9} \\ \frac{8}{9} & -\frac{1}{9} & \frac{4}{9} \\ \frac{4}{9} & \frac{4}{9} & -\frac{7}{9} \end{array} \right) . \tag{11}$$

A diagonal charged lepton mass matrix is invariant under an infinite choice of abelian symmetries since charge assigments for the left handed fields may be compensated by the corresponding right ones. A natural choice is given by $Z_e \times Z_\mu \times Z_\tau$. Clearly this symmetry has to be broken by soft terms or by NLO contributions if the correction to the TP mixing matrix has to arise by the charged lepton sector.

3 The model

In this section we build a renormalizable model that provides the TP mixing matrix. We assume that no other heavy matter fields exist a part from those we report in tab. 1, thus the Yukawa lagrangian we write in eq. (12) is complete and no NLO terms have to be taken into account. The model is based on the flavor symmetry $G_f \sim SU(3)_F \times U(1)_F$ and matter and scalar fields charge assigments are reported in tab.1. Left handed doublets transform as a triplet of $SU(3)_F$. Standard model (SM) right handed charged lepton are $SU(3)_F$ singlet and charged under $U(1)_F$. Among the matter fields we have 2 right handed neutrinos, singlet under $SU(3)_F \times U(1)_F$, a vectorial couple of heavy SM singlets Σ,$\overline{\Sigma}$, transforming as 3 and $\overline{3}$ respectively under $SU(3)_F$, another vectorial couple of heavy SM $SU(2)$ singlet charged under $U(1)_Y$, F, F^c, 3 and $\overline{3}$ of $SU(3)_F$ respectively. We introduce five $\overline{3}$ scalar fields, three of them are charged under $U(1)_F$. In addition we impose an extra Z_2 symmetry under which all the fields are even with the exception of one right handed neutrino, ν_2^c, and one scalar triplet, ϕ_2, that are odd.

Matter	L	ν_1^c	ν_2^c	e^c	μ^c	τ^c	Σ	$\bar{\Sigma}$	F	F^c	ϕ_1	ϕ_2	ϕ_e	ϕ_μ	ϕ_τ	H
$SU(2)_L$	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2
$U(1)_Y$	-1/2	0	0	1	1	1	0	0	-1	1	0	0	0	0	0	1/2
$SU(3)_F$	3	1	1	1	1	1	3	$\overline{3}$	3	$\overline{3}$	$\overline{3}$	$\overline{3}$	$\overline{3}$	$\overline{3}$	$\overline{3}$	1
$U(1)_F$	0	0	0	α	β	γ	0	0	0	0	0	0	$-\alpha$	$-\beta$	$-\gamma$	0

Table 1: *Transformation properties of the matter fields. The choice of the charged lepton* $U(1)_F$ *charges is arbitrary once the condition* $\alpha \neq \beta \neq \gamma$ *is satisfied.*

3.1 Mass matrices

Given the field content of tab.1 the lagrangian reads as

$$\begin{aligned}\mathcal{L} &= kLH\overline{\Sigma}+(y_1\Sigma\phi_1+y_1'\overline{\Sigma}\phi_1^*)\nu_1^c+(y_2\Sigma\phi_2+y_2'\overline{\Sigma}\phi_2^*)\nu^c{}_2+M_\Sigma\Sigma\overline{\Sigma}+\frac{M_1}{2}\nu_1^c\nu_1^c+\frac{M_2}{2}\nu_2^c\nu_2^c\\ &+ y_F L\tilde{H}F^c+y_eF\phi_e e^c+y_\mu F\phi_\mu\mu^c+y_\tau F\phi_\tau\tau^c+M_FFF^c\,, \end{aligned}\tag{12}$$

where $\tilde{H}=i\sigma_2 H$ with H the usual SM higgs doublet and σ_2 the Pauli matrix. We have omitted $SU(2)$ indices to simplify the notation. As already stated this is the full Yukawa lagrangian thus no NLO corrections have to been included.

In sec.2 we have said that TP mixing matrix is obtained when neutrino and charged lepton mass matrices are invariant under $Z_2\times Z_2$ and $Z_e\times Z_\mu\times Z_\tau$ respectively. This means that the flavor group G_f has to be broken following different patterns in the neutrino and charged lepton sector as it usually happens in the context of discrete flavour symmetries [5]. In our scenario this is realized when the five scalars ϕ_i develop vacuum expectation values (VEVs) as

$$\begin{aligned}&\langle\phi_1\rangle\sim(2,2,1)\,, &&\langle\phi_2\rangle\sim(2,-1,-2)\,,\\ &\langle\phi_e\rangle\sim(1,0,0)\,, &&\langle\phi_\mu\rangle\sim(0,1,0)\,,\quad\langle\phi_\mu\rangle\sim(0,0,1)\,.\end{aligned}\tag{13}$$

In sec.3.2 we sketch how this alignment may be realized.

When flavor and electroweak symmetries are broken the neutrino mass matrix is a 5 block matrix that in the basis $(\nu_L,\overline{\Sigma},\Sigma,\nu_1^c,\nu_2^c)$ is given by

$$M_\nu = \begin{pmatrix}0 & m_D & 0\\ m_D^T & M_0 & \lambda\\ 0 & \lambda & M_{\nu^c}\end{pmatrix}\,.\tag{14}$$

with

$$\begin{aligned}&m_D=(kv_h\cdot\mathbb{1},0)\,, &&\lambda=\begin{pmatrix}\lambda_1' & \lambda_2'\\ \lambda_1 & \lambda_2\end{pmatrix}\,,\\ &M_{\nu^c}=Diag(M_1,M_2)\,, &&M_0=\begin{pmatrix}0 & M_\Sigma\cdot\mathbb{1}\\ M_\Sigma\cdot\mathbb{1} & 0\end{pmatrix}\,,\end{aligned}\tag{15}$$

being $\mathbb{1}$ the identity 3×3 matrix. The λ's are defined as

$$\lambda_1,\lambda_1'=y_1,y_1'v_1\begin{pmatrix}2\\2\\1\end{pmatrix}\qquad\lambda_2,\lambda_2'=y_2,y_2'v_2\begin{pmatrix}2\\-1\\-2\end{pmatrix}\tag{16}$$

Under the assumption $M_\Sigma > M_{1,2} > \lambda_{1,2},\lambda_{1,2}'$ it is convenient defining the spinor Σ_1 and Σ_2

$$\Sigma_1=\frac{1}{\sqrt{2}}(\overline{\Sigma}+\Sigma)\quad\Sigma_2=\frac{1}{\sqrt{2}}(-\overline{\Sigma}+\Sigma)\,.\tag{17}$$

In this way eq. (14) becomes

$$M_\nu \;=\; \left(\begin{array}{ccc} 0 & \tilde{m}_D & 0 \\ \tilde{m}_D^T & \tilde{M}_0 & \tilde{\lambda} \\ 0 & \tilde{\lambda}^T & M_{\nu^c} \end{array}\right) . \tag{18}$$

M_ν may be sequentially diagonalized by using the block diagonalization method introduced in [16]. First the method is applied to the block involving the heavy fields $(\Sigma_1, \Sigma_2, \nu_1^c, \nu_2^c)$. The unitarity matrix that diagonalize the block is defined as

$$U_H =\sim \left(\begin{array}{cc} 1-\frac{BB^T}{2} & B \\ -B^T & 1-\frac{B^TB}{2} \end{array}\right) \tag{19}$$

with

$$B \sim -\tilde{M}_0^{-1}\tilde{\lambda} . \tag{20}$$

The lightest singlet neutrinos mass matrix becomes

$$\tilde{M}_{\nu^c} = M_{\nu^c} - \tilde{\lambda}^T \tilde{M}_0^{-1} \tilde{\lambda} . \tag{21}$$

The effective light neutrino mass matrix is given by the usual type I see saw formula according to

$$m_\nu \;\sim\; -\tilde{m}_D B \frac{1}{\tilde{M}_{\nu^c}} B^T m_D^T \sim -\tilde{m}_D \frac{1}{\tilde{M}_0}\tilde{\lambda}\frac{1}{M_{\nu^c}}\tilde{\lambda}^T \frac{1}{\tilde{M}_0^T} m_D^T , \tag{22}$$

and presents the form

$$m_\nu = \left(\begin{array}{ccc} 4x+4y & 4x-2y & 2x-4y \\ 4x-2y & 4x+y & 2x+2y \\ 2x-4y & 2x+2y & x+4y \end{array}\right) \tag{23}$$

with

$$x = 2k^2 v_h^2 \frac{y_1^2 v_1^2}{M_1 M_\Sigma^2} , \quad y = 2k^2 v_h^2 \frac{y_2^2 v_2^2}{M_2 M_\Sigma^2} . \tag{24}$$

The previous m_ν is diagonalized by U_{TP} with eigenvalues $(9x, 9y, 0)$. Thus our realization allow only the IH spectrum.

For what concerns the charged lepton sector in addition to the SM fields we have the heavy fields F, F^c. When the vacuum alignment coincides exactly with that in eq. (13) the full charged lepton left-right mass matrix presents a trivial block structure

$$M_{ch} = \left(\begin{array}{cc} 0 & y_F v_H \cdot \mathbb{1} \\ Y & M_F \cdot \mathbb{1} \end{array}\right) , \tag{25}$$

with $Y = \mathrm{Diag}(y_e v_e, y_\mu v_\mu, y_\tau v_\tau)$. By integrating out the heavy fields the SM charged lepton mass matrix is diagonal and

$$(m_e, m_\mu, m_\tau) = \frac{y_F v_H}{M_F}(y_e v_e, y_\mu v_\mu, y_\tau v_\tau) . \tag{26}$$

In sec.3.2 it is discussed how the alignments given in eq. (13) get corrections due to the presence of soft terms needed to give mass to the Goldstone Bosons (GBs) arising by minimizing the potential. Specifically, for what concerns $\phi_{e,\mu,\tau}$, eq. (13) have to be substituted by

$$\langle \phi_{e,\mu,\tau} \rangle + \epsilon_{e,\mu,\tau}(2,2,1)\,. \tag{27}$$

In this way the block Y is substituted by

$$\tilde{Y} = \begin{pmatrix} y_e(v_e + 2\epsilon_e) & 2y_\mu\epsilon_\mu & 2y_\tau\epsilon_\tau \\ 2y_e\epsilon_e & y_\mu(v_\mu + 2\epsilon_\mu) & 2y_\tau\epsilon_\tau \\ y_e\epsilon_e & y_\mu\epsilon_\mu & y_\tau(v_\tau + 2\epsilon_\tau) \end{pmatrix} \simeq \begin{pmatrix} y_e v_e & 2y_\mu\epsilon_\mu & 2y_\tau\epsilon_\tau \\ 2y_e\epsilon_e & y_\mu v_\mu & 2y_\tau\epsilon_\tau \\ y_e\epsilon_e & y_\mu\epsilon_\mu & y_\tau v_\tau \end{pmatrix}\,, \tag{28}$$

since $\epsilon_x << v_x$. Neglecting terms proportional to $y_e v_e << y_\mu v_\mu, y_\tau v_\tau$ the charged lepton mass matrix squared presents the following structure

$$m_{ch}m_{ch}^\dagger \simeq |m_\tau|^2 \begin{pmatrix} 0 & \epsilon'_\mu \frac{|y'_\mu|^2}{|y'_\tau|^2} & \epsilon'_\tau \\ \epsilon'^*_\mu \frac{|y'_\mu|^2}{|y'_\tau|^2} & \frac{|y'_\mu|^2}{|y'_\tau|^2} & \epsilon'_\tau + \epsilon'^*_\mu \frac{|y'_\mu|^2}{|'y_\tau|^2} \\ \epsilon'^*_\tau & \epsilon'^*_\tau + \epsilon'_\mu \frac{|y'_\mu|^2}{|y'_\tau|^2} & 1 \end{pmatrix}\,, \tag{29}$$

where we have defined $y'_x = y_x v_x, \epsilon'_x = y_x v_x \epsilon_x/|m_\tau|^2$. For construction $\epsilon'_x << 1$ and $|\epsilon'_\tau| < |\epsilon'_\mu|$ to fit the correct ratio $|m_\mu|/|m_\tau|$ thus eq. (29) is diagonalized by

$$U_e = \begin{pmatrix} 1 & \epsilon'_\mu & \epsilon'_\tau \\ \epsilon'^*_\mu & 1 & \epsilon'_\tau \\ \epsilon'^*_\tau & \epsilon'^*_\tau & 1 \end{pmatrix}\,. \tag{30}$$

The final lepton mixing matrix has the desired structure and in first approximation it is given by

$$U_{lep} = U_e^\dagger U_{TP} \simeq \begin{pmatrix} \frac{2}{3} - \frac{2\epsilon'_\mu}{3} & -\frac{\epsilon'_\mu}{3} - \frac{2}{3} & \frac{2\epsilon'_\mu}{3} + \frac{1}{3} \\ \frac{2\epsilon'^*_\mu}{3} + \frac{2}{3} & \frac{1}{3} - \frac{2\epsilon'^*_\mu}{3} & \frac{\epsilon'^*_\mu}{3} - \frac{2}{3} \\ \frac{1}{3} & \frac{2}{3} & \frac{2}{3} \end{pmatrix}\,, \tag{31}$$

where we have used $|\epsilon'_\tau| < |\epsilon'_\mu|$.

3.2 Vacuum alignment

Model based on flavor symmetries spontaneously broken in different directions in the charged lepton and neutrino sectors have always to face off the problem of realizing the correct vacuum alignments. This affects TBM as well as BM models based on both discrete and continuous symmetries. The formers tend to break in one direction and therefore different techniques have been developed to break them in two directions. For the latter the situation is even worse since continuous symmetry do not develop a preferred direction to be broken to and the minimum conditions present an infinite degeneracy. For this reason in model based on flavor continuous groups such as $SU(3)$ or $SO(3)$ the correct vacuum alignment is typically obtained by introducing soft breaking terms of the continuous symmetry. These softs preserve an appropriate discrete subgroup of the continuos symmetry and through the minimization of the potential they select the correct directions [17]. Here we use the same approach.

In our model we have five $\overline{3}$ of the flavour group $G_f \sim SU(3)_F \times U(1)_F \times Z_2$. For five $\overline{3}$ of a generic $SU(3)$ the most generic potential is written as

$$V[\phi_i] = \mu^2_{ij}\phi^\dagger_i\phi_j + \lambda_{ijkl}(\phi^\dagger_i\phi_j)(\phi^\dagger_k\phi_l)\,, \tag{32}$$

where the sum other all the fields is understood. In our case, the abelian symmetries forbid many terms and in particular the scalar $SU(3)_F$ invariant potential has an accidental enlarged symmetry $SU(3) \times U(3)^3$: the first $SU(3)$ involves ϕ_1 and ϕ_2 and one $U(3)$ per ϕ_i, $i = e, \mu, \tau$. The generic vacuum configuration constrains the absolute scalar VEVs

$$\langle\phi_1\rangle\ ,\quad \langle\phi_2\rangle\ ,\quad \langle\phi_{e,\mu,\tau}\rangle\ , \tag{33}$$

and the accidental $SU(3) \times U(3)^3$ is completely broken giving rise to $8 + 9 \times 3 = 35$ GBs. This is a situation quite common in flavor models: the inclusion of the soft terms is needed both to trigger the correct alignments and to give mass to the unwanted GBs.

In our context we need three set of soft terms:

- V^1_{soft}: it triggers the correct vacuum alignment for $\phi_{e,\mu,\tau}$ and also breaks the accidental $U(3)^3$ symmetry to $SU(3)$, giving mass to 19 GBs. At this level we are left with an accidental global symmetry $SU(3) \times SU(3)$
- V^2_{soft}: it triggers the correct vacuum alignment for $\phi_{1,2}$ breaking $SU(3) \times SU(3)$ to $SU(3)$ and giving mass to other 8 GBs.
- V^3_{soft}: it breaks the residual $SU(3)$ and gives mass to the last 8 GBs.

A suitable example for V^1_{soft} is given by

$$V^1_{soft} = [m^2_{e\mu}(\phi^\dagger_e\phi_\mu) + m^2_{\mu\tau}(\phi^\dagger_\mu\phi_\tau) + A\phi_e\phi_\mu\phi_\tau] + H.c. \tag{34}$$

The basic assumption is that the soft terms slightly modify the first derivative system that in first approximation may be considered unchanged. Thus the first derivative system of the potential involving $\phi_{e,\mu,\tau}$ fixes $\langle\phi_{e,\mu,\tau}\rangle$ and we have always the freedom to choose one direction, for example

$$\langle\phi_e\rangle = v_e(1, 0, 0)\,. \tag{35}$$

If $m^2_{e\mu}, m^2_{\mu\tau} > 0$ and $A < 0$ the quadratic and quartic terms of eq. (34) select orthogonal directions for ϕ_μ and ϕ_τ:

- the quadratic terms select the direction

$$\langle\phi_\mu\rangle = v_\mu(0, \cos\alpha, \sin\alpha), \langle\phi_\tau\rangle = v_\tau(0, \sin\alpha, -\cos\alpha)\,, \tag{36}$$

- the trilinear term is proportional to $\cos 2\alpha$ and to maximize it we need $\alpha = \pi/2$ that gives the correct alignment to ϕ_μ and ϕ_τ.

Building V^2_{soft} is a bit more *ad hoc*: we need to impose that V^2_{soft} is invariant under one of the following transformations

$$\begin{array}{ll} i) & \phi_{1_1} \to 2\phi_{1_3} \quad \phi_{1_3} \to 1/2\phi_{1_1}\,, \\ ii) & \phi_{1_1} \to \phi_{1_2} \quad \phi_{1_2} \to \phi_{1_1}\,, \\ iii) & \phi_1 \to \phi_2 \end{array} \tag{37}$$

Clearly this transformations breaks explicitly $SU(3)$ and the Z_2 under which ϕ_2 is odd. A possible V^2_{soft} is given by

$$m_1^2|\phi_{1_1} - 2\phi_{1_3}|^2 + m_\pm^2|\phi_{1_1} \pm \phi_{1_2}|^2 + m_{12}^2(\phi_1^\dagger\phi_2 + H.c.)\,, \tag{38}$$

By choosing correctly the sign of $m_1^2, m_-^2, m_{12}^2 > 0, m_+^2 < 0$ not to have tachyons we get $\langle\phi_1\rangle$ and $\langle\phi_2\rangle$ orthogonal and along the right directions.

Finally V^3_{soft} have to make massive the last 8 GBs: in order not to destroy the alignments provided by V^1_{soft} and V^2_{soft} it has to be subdominant with respect to them. It may have a form like

$$m_{\tau 1}^2(\phi_\tau^\dagger\phi_1)\,. \tag{39}$$

In general V^3_{soft} leaves the freedom to preserve only one vev direction and would slightly disalign the others. If only ϕ_1–or ϕ_2–is involved in V^3_{soft} together with $\phi_{e,\mu,\tau}$ it is possible to disalign only the triplets entering in the charged lepton Yukawa lagrangian. For example they would disalign according to

$$\langle\phi_{e,\mu,\tau}\rangle + \epsilon_{e,\mu,\tau}(2,2,1)\,, \tag{40}$$

giving rise to the corrections needed to generate a non trivial charge lepton mixing.

4 Neutrino phenomenological analysis

Eq.(23) and eq. (31) give us all the informations to outline the neutrino phenomenology of the model discussed. For what concerns the spectrum only the IH case is allowed with vanishing m_3. In first approximation the predictions for the lepton angles coincide with those given in eq. (6) but a more accurate scan of the parameters space is performed by taking into account the complete charged lepton mixing matrix obtained fitting the charged lepton masses. The result is showed in fig.3. From the upper to the lower panel we plot the reactor angle versus the solar angle, the atmospheric angle and the CP dirac phase δ_l. The numerical scan confirms the parametric plot showed in the introduction. We have reported the 3σ range for the 3 angles according to the most recent analysis. There is a nice correlation between the solar and reactor angle: if in the near future there would be an improvement in the 3σ range precision of one of the two angles we would automatically get an upper or/and lower predictions for the other. On the other hand the model could be ruled out by an improvement on the precision for the atmospheric angle since it predicts $\sin\theta_{23}^2$ far from its central value (0.42). For what concerns the CP Dirac phase our points clustered in the range $0 \pm \pi/4$. From eq. (7) we see that $\delta_l \sim 0$ is the expected value from the analytical parametrization. Indeed eq. (6) tell us that $\cos\delta \sim -1$ to fit the solar angle.

To study neutrinoless double beta decay we consider the effective $0\nu\beta\beta$ parameter m_{ee} defined as

$$m_{ee} = [U_{lep}\,\mathrm{diag}(m_1,\, m_2,\, 0)\, U_{lep}]_{11}. \tag{41}$$

In fig.4 we plot m_{ee} versus the $\sin\theta_{13}^2$ since in our model the lightest neutrino mass is always vanishing. As consequence the model predicts almost an exact value for $m_{ee} \sim 45$ meV. The future experiments are expected to reach good sensitivities: 90 meV [18] (GERDA), 20 meV [19] (Majorana), 50 meV [20] (SuperNEMO), 15 meV [21] (CUORE) and 24 meV [22] (EXO). As a result the model may be tested in the next future.

5 Conclusions

In this paper we have proposed a new mixing matrix for the neutrino mass matrix that we named the Tri-Permuting (TP) mixing matrix. This pattern requires large corrections both to the solar and to the reactor angles that thus are correlated in a new way orthogonal to other patterns proposed in literature such as the BM mixing one. Not to affect neutrino masses and the atmospheric angle these corrections have to arise by the charged lepton mixing matrix. We have build a full renormalizable model in which this scenario is realized. In the model proposed both neutrino and charged lepton get mass through a generalized see saw. The model is characterized by a neutrino IH spectrum with vanishing m_3. As consequence it is highly testable in the next future because it predicts an exact value for the effective $0\nu\beta\beta$ parameter $m_{ee} \sim 45$ meV and could be ruled out by an improvement of precision for the atmospheric angle. At the same time it gives a nice correlation between solar and reactor angles that could be tested by future analysis.

We have also roughly discussed the potential sketching the strategy to obtain the correct vacuum alignments and the correction to the charged lepton mass matrix needed to correct the TP mixing matrix. A part from the neutrino sector the model phenomenology is deeply rich due to the presence of many new scalars and heavy fermions. A complete analysis of its phenomenology is postponed to a future work [23].

Aknowledgments

I am grateful to S. Morisi for useful discussions and suggestions in the initial stage of this project.

References

[1] K. Abe *et al.* [T2K Collaboration], arXiv:1106.1238 [Unknown].

[2] MINOS Collaboration, http://www-numi.fnal.gov/pr_plots/index.html.

[3] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, arXiv:1106.6028 [hep-ph].

[4] T. Schwetz, M. Tortola and J. W. F. Valle, arXiv:1108.1376 [hep-ph].

[5] A complete list of reference may be found in G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701 (2010) [arXiv:1002.0211 [hep-ph]].

[6] S. F. King and C. Luhn, arXiv:1107.5332 [hep-ph]. X. G. He and A. Zee, arXiv:1106.4359 [hep-ph]. T. Araki, [arXiv:1106.5211 [hep-ph] S. Morisi, K. M. Patel and E. Peinado, arXiv:1107.0696 [hep-ph]. Y. H. Ahn, H. Y. Cheng and S. Oh, arXiv:1107.4549 [hep-ph].

[7] D. Marzocca, S. T. Petcov, A. Romanino, M. Spinrath, [arXiv:1108.0614 [hep-ph]].

[8] Z. z. Xing, arXiv:1106.3244 [hep-ph]. H. Zhang and S. Zhou, arXiv:1107.1097 [hep-ph]. W. Chao and Y. j. Zheng, arXiv:1107.0738 [hep-ph]. J. M. Chen, B. Wang and X. Q. Li, arXiv:1106.3133 [hep-ph]. N. Qin and B. Q. Ma, arXiv:1106.3284 [hep-ph]. J. E. Kim and M. S. Seo, arXiv:1106.6117 [hep- ph]. Y. j. Zheng and B. Q. Ma, arXiv:1106.4040 [hep-ph].

[9] V. D. Barger, S. Pakvasa, T. J. Weiler and K. Whisnant, Phys. Lett. B **437**, 107 (1998) [arXiv:hep-ph/9806387]; Y. Nomura and T. Yanagida, Phys. Rev. D **59** (1999) 017303 [arXiv:hep-ph/9807325]; G. Altarelli and F. Feruglio, Phys. Lett. B **439** (1998) 112 [arXiv:hep-ph/9807353].

[10] G. Altarelli, F. Feruglio and L. Merlo, JHEP **0905**, 020 (2009) [arXiv:0903.1940 [hep-ph]].

[11] D. Meloni, [arXiv:1107.0221 [hep-ph]].

[12] P. H. Frampton, S. T. Petcov and W. Rodejohann, Nucl. Phys. B **687** (2004) 31 [arXiv:hep-ph/0401206]; G. Altarelli, F. Feruglio and I. Masina, Nucl. Phys. B **689** (2004) 157 [arXiv:hep-ph/0402155]; M. Raidal, Phys. Rev. Lett. **93** (2004) 161801 [arXiv:hep-ph/0404046]; H. Minakata and A. Y. Smirnov, Phys. Rev. D **70** (2004) 073009 [arXiv:hep-ph/0405088]; P. H. Frampton and R. N. Mohapatra, JHEP **0501**, 025 (2005), hep-ph/0407139; J. Ferrandis and S. Pakvasa, Phys. Rev. D **71** (2005) 033004 [arXiv:hep-ph/0412038]; S. K. Kang, C. S. Kim and J. Lee, arXiv:hep-ph/0501029; N. Li and B. Q. Ma, arXiv:hep-ph/0501226; K. Cheung, S. K. Kang, C. S. Kim and J. Lee, arXiv:hep-ph/0503122; Z. z. Xing, arXiv:hep-ph/0503200; A. Datta, L. Everett and P. Ramond, arXiv:hep-ph/0503222; S. Antusch, S. F. King and R. N. Mohapatra, arXiv:hep-ph/0504007; M. Lindner, M. A. Schmidt and A. Y. Smirnov, arXiv:hep-ph/0505067; H. Minakata, arXiv:hep-ph/0505262; T. Ohlsson, arXiv:hep-ph/0506094; A. Dighe, S. Goswami and P. Roy, Phys. Rev. D **73** (2006) 071301 [arXiv:hep-ph/0602062]; B. C. Chauhan, M. Picariello, J. Pulido and E. Torrente-Lujan, Eur. Phys. J. C **50** (2007) 573 [arXiv:hep-ph/0605032]; K. A. Hochmuth and W. Rodejohann, Phys. Rev. D **75** (2007) 073001 [arXiv:hep-ph/0607103]; M. A. Schmidt and A. Y. Smirnov, Phys. Rev. D **74** (2006) 113003 [arXiv:hep-ph/0607232]; F. Plentinger, G. Seidl and W. Winter, Nucl. Phys. B **791** (2008) 60 [arXiv:hep-ph/0612169]; F. Plentinger, G. Seidl and W. Winter, Phys. Rev. D **76** (2007) 113003 [arXiv:0707.2379 [hep-ph]].

[13] R. d. A. Toorop, F. Feruglio and C. Hagedorn, arXiv:1107.3486 [hep-ph].

[14] F. Feruglio and A. Paris, JHEP **1103**, 101 (2011) [arXiv:1101.0393 [hep-ph]].

[15] S. F. King, Phys. Lett. **B 439** (1998) 350 [arXiv:hep-ph/9806440]; S. F. King, Nucl. Phys.**B 562** (1999) 57 [arXiv:hep-ph/9904210]; S. F. King, Nucl. Phys. **B 576** (2000) 85 [arXiv:hep-ph/9912492]; S. F. King, JHEP **0209** (2002) 011 [arXiv:hep-ph/0204360]; S. F. King, Phys. Rev. D **67** (2003) 113010 [arXiv:hep-ph/0211228].

[16] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).

[17] I. de Medeiros Varzielas, G. G. Ross, Nucl. Phys. **B733**, 31-47 (2006). [hep-ph/0507176].

I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **644**, 153 (2007) [arXiv:hep-ph/0512313].

[18] A. A. Smolnikov and f. t. G. Collaboration, arXiv:0812.4194 [nucl-ex].

[19] Majorana Collaboration, arXiv:0811.2446 [nucl-ex].

[20] H. Ohsumi [NEMO and SuperNEMO Collaborations], J. Phys. Conf. Ser. **120** (2008) 052054.

[21] A. Giuliani [CUORE Collaboration], J. Phys. Conf. Ser. **120** (2008) 052051.

[22] M. Danilov *et al.*, Phys. Lett. B **480** (2000) 12 [arXiv:hep-ex/0002003].

[23] F. Bazzocchi, in preparation.



Figure 3: *The predictions for the lepton mixing parameters. From the upper to the lower panel we plot the reactor angle versus the solar angle, the atmospheric angle and the CP dirac phase* δ_l*. Vertical and horizontal lines bound the* 3σ *range for the corresponding angles.*



Figure 4: *The predictions for* m_{ee}*. Our IH spectrum is characterized by a vanishing* m_3*. As consequence we predict almost an exact value for* $m_{ee} \sim 45\ meV$ *that is in the precision range of the forthcoming neutrino experiments. Vertical lines bound the* 3σ *range for* $\sin\theta_{13}^2$.